

A DIAGONAL REPRESENTATION OF QUANTUM DENSITY MATRIX USING q -BOSON OSCILLATOR COHERENT STATES

R.Parthasarathy and R.Sridhar¹

The Institute of Mathematical Sciences
C.I.T. Campus, Tharamani, Chennai- 600113, INDIA

Abstract

A q -analogue of Sudarshan's diagonal representation of the Quantum Mechanical density matrix is obtained using q -boson coherent states. Earlier result of Mehta and Sudarshan on the self reproducing property of $\rho(z', z)$ is also generalized and a self-consistent self-reproducing kernel $\tilde{K}(z', z)$ is constructed.

¹e-mail addresses: sarathy@imsc.ernet.in ; sridhar@imsc.ernet.in

A diagonal representation of the quantum mechanical density matrix using standard bosonic oscillator coherent states has been given by Sudarshan [1]. One of the remarkable features of this representation is that for any normal ordered operator, its "average expectation value", becomes the same as that of the classical function for a probability distribution over the complex plane. So, the classical complex representation and the quantum mechanical density matrices are in one-to-one correspondence.

The emergence of quantum groups in the study of inverse scattering [2] resulted in q-Boson [3] and q-Fermion [4] oscillators and many usual quantum mechanical descriptions find their appropriate q-analogues [5]. It is possible that q-oscillators represent an effective way of dealing with non-ideal systems in which interactions are incorporated [6]. It is not obvious at present, whether nature makes use of q-Bosons (or q-Fermions) as some kind of non-linear excitations of the electromagnetic (or electron) fields. Nevertheless, it is worthwhile to examine the q-analogue of important quantum mechanical descriptions. It is the purpose here to find a q-analogue of the diagonal representation of the Quantum Mechanical density matrix. Although Nelson and Fields [7] have outlined a q-analogue of the diagonal representation of the density matrix, our results, while in agreement with them, offer further insight into the problem, such as the relationship between q-coherent state matrix elements and Fock space matrix elements, self-reproducing property and self reproducing kernel.

The q-Boson creation $a^\dagger$ and annihilation a operators satisfy

$$aa^\dagger - qa^\dagger a = 1, \quad ; \quad [N, a] = -a, \quad [N, a^\dagger] = a^\dagger, \quad (1)$$

where $N(\neq a^\dagger a)$ is the number operator. The q-Boson vacuum state $|0\rangle$ is an element of Fock Space $\mathcal{F}$ and $a|0\rangle = 0$; $\langle 0|0\rangle = 1$. The n q-Boson normalized state $|n\rangle$ is given by

$$|n\rangle = \frac{1}{\sqrt{[n]!}} (a^\dagger)^n |0\rangle, \quad (2)$$

where $\{|n\rangle | n \in \mathbf{N} \cup 0\}$ and the q-number $[n]$ is

$$[n] = \frac{1 - q^n}{1 - q} ; \quad 0 \leq q \leq 1, \quad (3)$$

such that $[n] \rightarrow n$ as $q \rightarrow 1$. The actions of $a, a^\dagger$ and N on $|n\rangle$ are

$$\begin{aligned} a |n\rangle &= \sqrt{[n]} |n-1\rangle, \\ a^\dagger |n\rangle &= \sqrt{[n+1]} |n+1\rangle, \quad N |n\rangle = n |n\rangle. \end{aligned} \quad (4)$$

and $[n]! = [n][n-1][n-2]\dots[1]$. The q-Boson coherent state [3,8] is given by

$$|z\rangle = \left(e_q^{|z|^2}\right)^{-\frac{1}{2}} \sum_{n=0}^{\infty} \frac{z^n}{\sqrt{[n]!}} |n\rangle, \quad (5)$$

where

$$e_q^x = \sum_{n=0}^{\infty} \frac{x^n}{[n]!}. \quad (6)$$

In the above $z \in C$ and e_q^x is the q-exponential function. Using (4) it readily follows that $|z\rangle$ is an eigen state of a with z as eigenvalue, i.e., $a |z\rangle = z |z\rangle$. If $|z'\rangle$ is another such state, then

$$\langle z | z' \rangle = \left(e_q^{|z|^2} e_q^{|z'|^2}\right)^{-\frac{1}{2}} e_q(\bar{z}z') \quad (7)$$

showing such states are *not* orthogonal. Nevertheless, $\langle z | z \rangle = 1$. Since $z \in C$ is continuous and $\langle z | z' \rangle$ is never vanishing, the states (5) satisfy the criteria of coherent states [9]. Furthermore, the states (5) provide a resolution of unity, namely, [8]

$$\frac{1}{\pi} \int d^2z |z\rangle \langle z| = 1. \quad (8)$$

Writing $z = re^{i\theta}$, $0 \leq \theta \leq 2\pi$, θ integration is ordinary Riemannian type while the r -integration will be "q-integration" [10], it follows from (5) that

$$e_q^{r^2} |re^{i\theta}\rangle \langle re^{i\theta}| = \sum_{n,m=0}^{\infty} r^{m+n} e^{i(n-m)\theta} \frac{1}{\sqrt{[n]![m]!}} |n\rangle \langle m|. \quad (9)$$

Multiplying this by $e^{i\ell\theta}$ and integrating over θ , we find

$$\int_0^{2\pi} \frac{d\theta}{2\pi} e_q^{r^2} e^{i\ell\theta} | r e^{i\theta} \rangle \langle r e^{i\theta} | = \sum_{n,m=0}^{\infty} \frac{r^{m+n}}{\sqrt{[n]![m]!}} | n \rangle \langle m | \delta(\ell + n - m). \quad (10)$$

Operating on (10) by the q -differential operator $(\frac{d}{dr})_q$ p -times, we find

$$\left(\frac{d}{dr}\right)_q^p \int_0^{2\pi} \frac{d\theta}{2\pi} e_q^{r^2} e^{i\ell\theta} | r e^{i\theta} \rangle \langle r e^{i\theta} | = \sum_{n,m=0}^{\infty} \frac{r^{m+n-p}}{\sqrt{[n]![m]!}} \frac{[m+n]!}{[m+n-p]!} | n \rangle \langle m | \delta(\ell + n - m). \quad (11)$$

Now, if we evaluate this equation at $r = 0$, it is easily seen that in the right hand side, the term with $m + n - p = 0$ alone survives. Thus

$$\left\{ \left(\frac{d}{dr}\right)_q^p \int_0^{2\pi} \frac{d\theta}{2\pi} e_q^{r^2} e^{i\ell\theta} | r e^{i\theta} \rangle \langle r e^{i\theta} | \right\}_{r=0} = \sum_{n,m=0}^{\infty} [m+n]! \frac{1}{[m+n-p]! \sqrt{[n]![m]!}} | n \rangle \langle m | \delta(\ell + n - m) \delta(n + m - p),$$

leading to the result

$$| n \rangle \langle m | = \frac{\sqrt{[n]![m]!}}{[n+m]!} \left(\frac{d}{dr}\right)_q^{(n+m)} \int_0^{2\pi} \frac{d\theta}{2\pi} e_q^{r^2} e^{i(m-n)\theta} | r e^{i\theta} \rangle \langle r e^{i\theta} | \Big|_{r=0}. \quad (12)$$

This result will be used later. A density matrix, in terms of the states in $\mathcal{F}$, has the following expansion:

$$\rho = \sum_{n,m=0}^{\infty} \rho(n, m) | n \rangle \langle m |. \quad (13)$$

Using (12) in (13), we have,

$$\rho = \sum_{n,m=0}^{\infty} \rho(n, m) \frac{\sqrt{[n]![m]!}}{[n+m]!} \left(\frac{d}{dr}\right)_q^{(n+m)} \int_0^{2\pi} \frac{d\theta}{2\pi} e_q^{r^2} e^{i(m-n)\theta} | r e^{i\theta} \rangle \langle r e^{i\theta} | \Big|_{r=0}. \quad (14)$$

which is the desired q-analogue of Sudarshan's diagonal representation of the density matrix.

Expansion of ρ in terms of the q-coherent states is

$$\rho = \int d^2 z \phi(z) |z\rangle\langle z| \quad (15)$$

where $d^2 z = dr^2 d\theta$, with dr^2 integration as q- integration. The complex function $\phi(z)$ is such that $\int d^2 z \phi(z) = 1$ due to $Tr(\rho) = 1$. The expansion coefficients (13) and (15) are related by

$$\rho(n, m) = \int d^2 z \phi(z) \frac{z^n \bar{z}^m}{\sqrt{[n]![m]!}} (e_q |z|^2)^{-1} \quad (16)$$

which is the q-analogue of Eqn.(7) of Sudarshan [1]. Performing the θ integration,

$$\rho(n, m) = \int_0^{R_q^2} \phi(r^2) \left\{ \frac{r^{2n}}{[n]!} e_q^{r^{2-1}} \right\} \delta_{n,m}, \quad (17)$$

consistent with $Tr\rho = 1$. It is to be noted that the q-exponential function necessitates the upper limit as $R_q^2 = \frac{1}{(1-q)}$, the first zero of the q-exponential function [7]. The expression in the curly bracket in (17) can be interpreted as the q-boson Poisson distribution[17].

As a first application of the q-coherent states, we consider the expansion of a bounded operator F as a normal ordered power series in a and $a^\dagger$, namely,

$$F = \sum_{p,q} C_{p,q} a^{\dagger p} a^q, \quad (18)$$

with $C_{p,q}$ as expansion coefficients. Using $a |z\rangle = z |z\rangle$, we have

$$\frac{\langle z' | F | z \rangle}{\langle z' | z \rangle} = \sum_{p,q} C_{p,q} \bar{z}'^p z^q. \quad (19)$$

On the other hand, using (5),

$$\frac{\langle z' | F | z \rangle}{\langle z' | z \rangle} = e_q^{\bar{z}' z^{-1}} \sum_{n,m} \frac{\langle n | F | m \rangle}{\sqrt{[n]![m]!}} (\bar{z}')^n z^m \quad (20)$$

From (19) and (20), we have

$$C_{p,q} = \sum_{k=0}^{Min(p,q)} \frac{(-1)^k q^{\frac{k(k-1)}{2}}}{[k]! s_q r t [q-k]! [p-k]!} \langle p-k | F | q-k \rangle. \quad (21)$$

$C_{p,q}$ in Eq(19) can be interpreted as the expansion coefficient of the operator F in the q-coherent state space, while $\langle n | F | m \rangle$, the corresponding coefficient in the Fock space and (21) relates them.

As a second application of the representation of ρ in terms of q-coherent states, we first consider the matrix element of ρ between such states. Using (5), (8) and (13), we find

$$\langle z' | \rho | z \rangle = \sum_{n,m} \rho(m,n) \left\{ e_q^{|z|^2} e_q^{|z'|^2} \right\}^{-\frac{1}{2}} \frac{\bar{z}'^m z^n}{\sqrt{[m]! [n]!}} \quad (22)$$

Introduce a function $\rho(z', z)$ as

$$\rho(z', z) = \sum_{m,n} \rho(m,n) \frac{\bar{z}'^m z^n}{\sqrt{[m]! [n]!}} \quad (23)$$

so that using (22)

$$\rho(z', z) = \sqrt{e_q^{|z|^2} e_q^{|z'|^2}} \langle z' | \rho | z \rangle. \quad (24)$$

The over completeness of the q-coherent states allows us to rewrite (24) as

$$\rho(z', z) = \frac{1}{\pi} \int d^2 \xi \langle z' | \rho | \xi \rangle \langle \xi | z \rangle \left\{ e_q^{|z|^2} e_q^{|z'|^2} \right\}^{\frac{1}{2}}. \quad (25)$$

Substituting for $\langle \xi | z \rangle$ from (7), (25) can be expressed as

$$\rho(z', z) = \int d^2 \xi K(z, \xi) \rho(z', \xi), \quad (26)$$

where

$$K(z, \xi) = \frac{1}{\pi} \{ e_q^{|\xi|^2} \}^{-1} e_q^{\bar{\xi} z}. \quad (27)$$

Eq(26) attributes the remarkable "self-reproducing property" for $\rho(z', z)$, with the kernel $K(z, \xi)$ in (27). Eqns (26) and (27) provide the q-analogue of eqns (3.25) and (3.26) of Mehta and Sudarshan[11].

From (27), it follows

$$\int d^2\xi K(z, \xi) K(\xi, z') = K(z, z'), \quad (28)$$

exhibiting the matrix structure of the kernels. The theory of reproducing kernels has been investigated by Aronszajan[12], according to which the hermitian nature of ρ demands the kernel $K(z, \xi)$ to satisfy

$$K(z, \xi)^* = K(\xi, z). \quad (29)$$

The above kernel (27), as well as the kernel (3.26) of Mehta and Sudarshan do not satisfy (29). This can be redressed by finding a suitable reproducing kernel. Instead of eq(23) for $\rho(z', z)$, we set

$$\tilde{\rho}(z', z) = \sum_{m,n} \rho(m, n) \frac{\bar{z}'^m z^n}{\sqrt{[m]![n]!}} \{e_q^{|z|^2} e_q^{|z'|^2}\}^{-\frac{1}{2}}, \quad (30)$$

which differs from (23) only by the q-exponentials. As these involve $|z|^2$ and $|z'|^2$, the analyticity of $\rho(z', z)$ and $\tilde{\rho}(z', z)$ are the same. With this definition, it follows that

$$\tilde{\rho}(z', z) = \int d^2\xi \tilde{K}(\xi, z) \tilde{\rho}(z', \xi), \quad (31)$$

with

$$\tilde{K}(\xi, z) = \frac{1}{\pi} \{e_q^{|\xi|^2} e_q^{|z|^2}\}^{\frac{-1}{2}} e_q^{\bar{\xi}z}. \quad (32)$$

This reproducing kernel has the desired reproducing property and satisfies eq(29), namely the hermiticity requirement. Furthermore, it is seen that

$$\tilde{K}(z, z) \geq 0, \quad (33)$$

and $\tilde{K}(z, z)$ is an entire function.

To summarize, we have constructed an explicit diagonal representation of the quantum mechanical density matrix using q-deformed boson coherent states. By expanding the density matrix in terms of q-boson coherent states ($\in C^\infty$) and then in terms of q-boson Fock space states ($\in \mathcal{F}$), the two expansion coefficients are shown to be related. The matrix element of ρ between

q-boson coherent states facilitated the introduction of $\rho(z', z)$ (23) which has the remarkable self-reproducing property (26) with respect to a kernel $K(z', z)$ in (27). This acquires significance due to the non-orthogonality of the q-boson coherent states (7). This provides a q-generalization of the earlier result of Mehta and Sudarshan [11]. A slightly modified reproducing kernel is introduced which satisfies all the required mathematical properties of a self-reproducing kernel for density matrix which is hermitian.

References

1. Sudarshan, E.C.G.: *Equivalence of semiclassical and quantum mechanical descriptions of statistical light beams*, Phys.Rev.Lett. **10** (1963) 277-279.
2. Drinfeld, V.G.: *Hopf algebras and the quantum Yang-Baxter Equation*, Soviet Math. Dokl **32** (1985) 254-258
Jimbo, M.: *A q-difference analogue of $U(g)$ and the Yang-Baxter equation*, Lett.Math.Phys. **10** (1985) 63-69.
3. Macfarlane, A.J.: *On q-analogues of the quantum harmonic oscillator and the quantum group $SU(2)_q$* , J.Phys.A: Math.Gen. **22** (1989) 4581-4588.
Biedenharn, L.C.: *The quantum group $SU_q(2)$ and a q-analogue of the boson operators*, J.Phys.A: Math.Gen. **22** (1989) L873-L878.
4. Parthasarathy, R and Viswanathan, K.S.: *A q-analogue of the supersymmetric oscillator and its q-super algebra*, J.Phys.A: Math.Gen. **24** (1991) 613 - 617.
5. Finkelstein, R.J.: *Gauged q-Fields*, hep-th/9906135; *Observable properties of q-deformed physical systems*, Lett.Math.Phys. **49** (1999) 105-114; *Spontaneous Symmetry Breaking of q-Gauge Field Theory*, hep-th/9908210; *TCP in q-Lorentz theories* Lett.Math.Phys. **54** (2000) 157-163; *On q-Electroweak*, hep-th/0110075; *On q-SU(3) Gauge Theory*, hep-th/0206067.

- Chaturvedi,S, Jagannathan,R, Srinivasan,V and Sridhar,R.: *Non-relativistic quantum mechanics in a non-commutative space*, J.Phys.A: Math.Gen. **26** (1993) L105-L112.
6. Esteve,J.G, Tejel,C and Villarroya,B.E.: *$SU_q(2)$ quantum group analysis of rotational spectra of diatomic molecules*, J.Chem.Phys. **96** (1992) 5614 - 5617;
 Chang,Z, Guo,H-Y and Yan,H.: *The q -deformed oscillator model and the vibrational spectra of diatomic molecules*, Phys.Lett. **A156** (1991) 192 - 196;
 Parthasarathy,R.: *q -Fermions*, IMSc-Preprint-93/23, April, 1993.
 7. Nelson,C.A and Fields,M.H.: *Number and Phase uncertainties of the q -analogue quantized field*. Phys.Rev. **A51** (1995) 2410 - 2429.
 8. Perelemov,A.M.: *On the completeness of some sub systems of q -deformed coherent states*, q-alg/9607006.
 9. Klauder,R.J and Skagerstam,B.: Eds *Coherent States: Applications in Physics and Mathematical Physics*, World Scientific (Singapore), 1985.
 10. Exton,M.: *q -Hypergeometric functions and applications*, Ellis Harwood Limited.
 11. Mehta,C.L and Sudarshan,E.C.G.: *Relation between quantum and semiclassical descriptions of optical coherence*, Phys.Rev. **B138** (1965) 274-280.
 12. Aronszajn,N.: *Theory of reproducing kernels*, Trans.Am.Math.Soc. **68** (1950) 337-404.